

EBIHARA, K.; NAKAJIMA, A.: Effect of Acetic Acid and Vinegar on Blood Glucose and Insulin Responses to Orally Administered Sucrose and Starch. Agric. Biol. Chem. 52 (1988) 1311 – 1312

FUSHIMI, T.; TAYAMA, K.; FUKAYA, M.; KITAKOSHI, K.; NAKAI, K., TSUKAMOTO, Y.; SATO, Y.: Acetic Acid Feeding Enhances Glycogen Repletion in Liver and Skeletal Muscle of Rats. The Journal of Nutrition 131 (2001) 1973 – 1977

FUSHIMI, T.; SATO, Y.: Effect of acetic acid feeding on the circadian changes in glycogen and metabolites of glucose and lipid in liver and skeletal muscle of rats. British Journal of Nutrition 94 (2005) 714 – 719

FUSHIMI, T.; SURUGA, K.; OSHIMA, Y.; FUKIHARU, M.; TSUKAMOTO, Y.; GODA, T.: Dietary acetic acid reduces serum cholesterol and triacylglycerols in rats fed a cholesterol-rich diet. British Journal of Nutrition 95 (2006) 916 – 924

JOHNSTON, C. S.; KIM, C. M.; BULLER, A. J.: Vinegar Improves Insulin Sensitivity to a High-Carbohydrate Meal in Subjects With Insulin Resistance or Type 2 Diabetes. Diabetes Care 27 (2004) 281 – 282

JOHNSTON, C. S.; BULLER, A. J.: Vinegar and Peanut Products as Complementary Foods to Reduce Postprandial Glycemia. Journal of the American Dietetic Association 105 (2005) 1939 – 1942

JOHNSTON, C. S.; GAAS, C. A.: Vinegar: Medicinal Uses and Antiglycemic Effect. MedGenMed. 8 (2006) 61

JOHNSTON, C. S.; WHITE, A. M.; KENT, S. M.: Preliminary evidence that regular vinegar ingestion favourably influences haemoglobin A1c values in individuals with type 2 diabetes mellitus. Diabetes Research and Clinical Practice 84 (2009) e15 – e17

JOHNSTON, C. S.; STEPLEWSKA, I.; LONG, C. A.; HARRIS, L. N.: Examination of the Antiglycemic Properties of Vinegar in Healthy Adults. Ann Nutr Metab 56 (2010) 74 – 79

KONDO, T.; KISHI, M.; FUSHIMI, T.; UGAJIN, S.; KAGA, T.: Vinegar Intake Reduces Body Weight, Body Fat Mass, and Serum Triglyceride Levels in Obese Japanese Subjects. Biosci. Biotechnol. Biochem. 73 (2009) 1837 – 1843

KONDO, T.; KISHI, M.; FUSHIMI, T.; KAGA, T.: Acetic Acid Upregulates the Expression of Genes for Fatty Acid Oxidation Enzymes in Liver To Suppress Body Fat Accumulation. J. Agric. Food Chem. 57 (2009) 5982 – 5986

LIATIS, S.; GRAMMATIKOU, S.; POULIA, K.; PERREA, D.; MAKRILAKIS, K.; DIAKOUMOPOULOU, E.; KATSILAMBROS, N.: Vinegar reduces postprandial hyperglycaemia in patients with type II diabetes when added to a high, but not to a low, glycaemic index meal. European Journal of Clinical Nutrition 64 (2010) 727 – 732

LILJEBERG, H.; BJÖRCK, I.: Delayed gastric emptying rate may explain improved glycaemia in healthy subjects to a starchy meal with added vinegar. European Journal of Clinical Nutrition 52 (1998) 368 – 371

LEEMAN, M.; ÖSTMAN, E.; BJÖRCK, I.: Vinegar dressing and cold storage of potatoes lowers postprandial glycaemic and insulinaemic responses in healthy subjects. European Journal of Clinical Nutrition 59 (2005) 1266 – 1271

METTLER, S.; SCHWARZ, I.; COLOMBANI, P. C.: Additive postprandial blood glucose-attenuating and satiety-enhancing effect of cinnamon and acetic acid. Nutrition Research 29 (2009) 723 – 727

MITROU, P.; RAPTIS, A.; LAMBADIARI, V.; BOUTATI, E.; PETSIOU, E.; SPANOUDI, F.; PAPAKONSTANTINOU, E.; MARATOU, E.; ECONOMOPOULOS, T.; DIMITRIADIS, G.; RAPTIS, S. A.: Vinegar Decreases Postprandial Hyperglycemia in Patients With Type 1 Diabetes. Diabetes Care 33 (2010) 27

MOHAMMADZADEH, S.; PROKHOROV, L. U.: Effect of Acetic Acid and Ethanol on Growth and Life Span of Chinese Hamster Tumor Cells. CANCER Bulletin 2 (2010) 53 – 59

NANDA, K.; MIYOSHI, N.; NAKAMURA, Y.; SHIMOJI, Y.; TAMURA, Y.; NISHIKAWA, Y.; UENAKAI, K.; KOHNO, H.; TANAKA, T.: Extract of Vinegar “Kurosu” from Unpolished Rice Inhibits the Proliferation of Human Cancer Cells. J. Exp. Clin. Cancer Res. 23 (2004) 69 – 75

OGAWA, N.; SATSU, H.; WATANABE, H.; FUKAYA, M.; TSUKAMOTO, Y.; MIYAMOTO, Y.; SHIMIZU, M.: Acetic Acid Suppresses the Increase in Disaccharidase Activity That Occurs during Culture of Caco-2 Cells. The Journal of Nutrition 130 (2000) 507 – 513

O’KEEFE, J. H.; GHEEWALA, N. M.; O’KEEFE, J. O.: Dietary Strategies for Improving Post-Prandial Glucose, Lipids, Inflammation, and Cardiovascular Health. Journal of the American College of Cardiology 51 (2008) 249 – 255

SAKAKIBARA, S.; YAMAUCHI, T.; OSHIMA, Y.; TSUKAMOTO, Y.; KADOWAKI, T.: Acetic acid activates hepatic AMPK and reduces hyperglycemia in diabetic KK-A(y) mice. Biochemical and Biophysical Research Communications 344 (2006) 597 – 604

SHIMOJI, Y.; SUGIE, S.; KOHNO, H.; TANAKA, T.; NANDA, K.; TAMURA, Y.; NISHIKAWA, Y.; HAYASHI, R.; UENAKAI, K.; OHIGASHI, H.: Extract of Vinegar “Kurosu” from Unpolished Rice Inhibits the Development of Colonic Aberrant Crypt Foci Induced by Azoxymethane. J. Exp. Clin. Cancer Res. 22 (2003) 591 – 597

SHIZUMA, T.; ISHIWATA, K.; NAGANO, M.; MORI, H.; FUKUYAMA, N.: Protective effects of fermented rice vinegar sediment (*Kurozu moromimatsu*) in a diethylnitrosamine-induced hepatocellular carcinoma animal model. J. Clin. Biochem. Nutr. 49 (2011) 31 – 35

SHIZUMA, T.; TANAKA, C.; MORI, H.; FUKUYAMA, N.: Protective Effects of Japanese Black Vinegar “*Kurozu*” and Its Sediment “*Kurozu Moromimatsu*” on Dextran Sulfate Sodium-induced Experimental Colitis. Dr. Fukata (Editor), InTech (2012)

WHITE, A. M.; JOHNSTON, C. S.: Vinegar Ingestion at Bedtime Moderates Waking Glucose Concentrations in Adults with Well-Controlled Type 2 Diabetes. *Diabetes Care* 30 (2007) 2814 – 2815